



Europäisches
Patentamt

European
Patent Office

Office européen
des brevets

PCT/EP 03/09302
10/524735

18 SEP 2003

21 AUG 2003

REC'D 29 SEP 2003

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02018772.0

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

BEST AVAILABLE COPY

R C van Dijk



Anmeldung Nr:
Application no.: 02018772.0
Demande no:

Anmeldetag:
Date of filing: 22.08.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Novartis Consumer Health S.A.
Case Postale 269
1260 Nyon
SUISSE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Topical composition

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K9/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

Topical Composition

The invention concerns topical formulations comprising the well-known and widely used non-steroidal anti-inflammatory drug (NSAID) diclofenac. It is one object of the present invention to provide an advantageous emulsion-gel specifically comprising the sodium salt of diclofenac, diclofenac Na. The latter represents the most simple, cheapest, most often used (taking into account oral and other routes of administration) and best studied derivative of diclofenac.

In US patent 4,917,886, a said emulsion-gel was disclosed already [see columns 2-3, composition B)]. However, the composition given there was an experimental one only and never commercialized. Its main purpose was to provide a direct comparison with the analogous crème composition A) comprising diclofenac Na and to demonstrate the superiority of emulsion-gels over ordinary crèmes. It had the following drawbacks: In order to obtain a sufficiently viscous emulsion-gel, especially in the presence of high amounts of lower alkanol (20% of isopropanol) and of an electrolyte (1% diclofenac Na), 1.2% of diethanolamine had to be used for neutralizing the gel-forming agent, polyacrylic acid (Carbopol 934 P).

However, nowadays the use of diethanolamine in topical formulations to be applied in humans is not allowed due to safety concerns of possible nitrosamine formation. It is therefore an object of the present invention to provide an emulsion-gel comprising diclofenac Na, which emulsion-gel can be obtained without neutralization with diethanolamine or other organic amines.

Moreover, a "true" emulsion-gel should be provided wherein diclofenac is fully dissolved rather than partly suspended, with identifiable crystals of diclofenac being present in the formulation. This is crucial to guarantee a high, constant and reproducible permeation of diclofenac through the skin.

These aims have been achieved, surprisingly, by selecting specific components in specific amounts and so being able to neutralize the composition with ammonia, NaOH or KOH, in particular ammonia.

The invention is further characterized in that the beneficial topical formulations provided do include diclofenac, and diclofenac Na specifically, in unusually low amounts only (<0.5%, w/w). Said low dose formulations are particularly advantageous in the topical treatment of burns including sunburn. They are providing similar relief and healing as normal dose formulations (e.g. 1%) but lead to much lower exposure of the body to the active substance.

Therefore the invention relates to a pharmaceutical composition intended for topical use, which composition is in the form of an opaque emulsion-gel and which composition comprises

- (a) 0.02-0.4% (w/w) of diclofenac sodium salt,
- (b) at least 50% (w/w) of water,
- (c) 0-30% (w/w) of at least one C2-C4-alkanol,
- (d) 3-20% (w/w) of a glycol solvent selected from the group consisting of propylene glycol and polyethylene glycol (200-1000),
- (e) 0.2-3% (w/w) of at least one gelling agent selected from the group consisting of carbomers,
- (f) 2-8% (w/w) of at least one lipid forming the oily phase of the emulsion-gel,
- (g) 1-5% (w/w) of at least one non-ionic surfactant, and
- (h) a basic agent selected from the group consisting of ammonia, sodium hydroxide and potassium hydroxide to adjust the pH of the total composition to 6.5-8.

All percentages given are percentages by weight (w/w), if not indicated otherwise.

Preferred are those compositions that comprise

- (a) 0.05-0.3% of diclofenac sodium salt,
- (b) 60-92% of water,
- (c) 0-25% of ethanol, isopropanol, or mixtures thereof,
- (d) 3-20% of propylene glycol,
- (e) 0.3-2% of at least one gelling agent selected from the group consisting of carbomers,
- (f) 3-7% of at least one lipid forming the fatty phase of the emulsion-gel,
- (g) 1-3% of at least one non-ionic surfactant, and
- (h) ammonia to adjust the pH of the total composition to 6.5-8.

In one embodiment of the invention, (c) is present in an amount of from 5 up to 25% and (d) is present in an amount of from 3 up to 7%, of the total composition. In another embodiment of the invention, (c) is present (or not, respectively) in an amount of from 0 up to 5%, in particular 0%, and (d) is present in an amount of from 3 up to 20% of the total composition.

In a further embodiment of the invention, (c) is present in an amount of from 0 up to 5% and (d) is present in an amount of from 3 up to 7% of the total composition. Said emulsion-gels have the additional advantage of being perceived by patients as extraordinarily pleasant and non-sticky on the skin in consumer tests. Said advantage is especially pronounced, if said compositions, in addition, comprise as component (f) isopropyl myristate, and in particular a mixture of isopropyl myristate and C6-C12-alkanoic acid C12-C18-alkyl esters.

Preferably, diclofenac Na (a) is present in amount of from 0.05 up to 0.3% - especially of from 0.1 up to 0.25% - of the total composition.

Water (b) is preferably present in an amount of from 50 up to 92%, in particular of from 60 up to 90%, or of from 60 up to 80%, of the total composition.

Preferred as C2-C4-alkanols (c) are ethanol, isopropanol, or mixtures thereof; and in particular isopropanol. Preferably, (c) is present in an amount of from 0 up to 25%, in particular of from 0 up to 20%, of the total composition. In one embodiment, (c) is typically present in an amount of from 5 up to 25%, and in particular of from 10 up to 20%, of the total composition. In another embodiment of the invention, (c) is present (or not, respectively) in an amount of from 0 up to 5% - preferably of from 0 up to 4%, more preferably of from 0 up to 3%, especially of from 0 up to 2%, and more especially of from 0 up to 1% - of the total composition.

Preferably, the glycol solvent (d) is propylene glycol. Propylene glycol is 1,2-propanediol. The glycol solvent (d) is preferably present in an amount of from 3.5 up to 20%, and in particular of from 4 up to 18%, of the total composition. In one embodiment, (d) is typically present in an amount of from 3 up to 7%, and in particular of from 3.5 up to 6%, of the total composition. In another embodiment of the invention, (d) is present in an amount of from 3 up to 20%, especially of from 4 up to 18%, of the total composition.

(e) Carbomers, in the context of the present invention, are defined as homo- or copolymers of acrylic acid that are cross-linked, e.g. with an allyl ether of pentaerythritol (allyl pentaerythritol) or an allyl ether of sucrose (allyl sucrose). Copolymers are formed e.g. with minor levels of long chain alkyl acrylate co-monomers. Homopolymers are preferred. Especially preferred are carbomers 980, 940, 981, 941, 974, 934 and 910. In particular preferred are the following products provided by Noveon, Inc, Cleveland, Ohio, USA (formerly B F Goodrich): Carbopol® 980 and Carbopol® 974, and analogous carbomer products from other suppliers. Preferably, (e) is present in an amount of from 0.3 up to 2%.

(f) Lipids form the oily phase of the emulsion-gels of the invention. They can be of a vegetable or animal nature, or partly or completely synthetic. There come into consideration lipids without ester linkages, e.g. hydrocarbons, fatty alcohols or fatty acids, and lipids having ester linkages, e.g. glycerides - i.e. fatty acid esters of glycerol -, or esters of fatty acids with C1-C36-alkanols. Hydrocarbons are e.g. paraffin or petroleum jelly. Fatty alcohols are e.g. decanol, dodecanol, tetradecanol, hexadecanol or octadecanol. Fatty acids are e.g. C6-C24-, especially C10-C18-, alkanolic acids, e.g. hexanoic acid, octanoic acid, decanoic acid, dodecanoic acid, tetradecanoic acid, hexadecanoic acid or octadecanoic acid. Fatty acids further include unsaturated fatty acids, e.g. oleic acid or linoleic acid. Glycerides are e.g. olive oil, castor oil, sesame oil, it being possible for all said oils also to be hydrogenated; caprylic/capric acid triglyceride or glycerol mono-, di- and tri-esters with palmitic and/or stearic acid. Esters of fatty acids with C1-C36-alkanols are e.g. beeswax, carnauba wax, cetyl palmitate, lanolin, isopropyl myristate, isopropyl stearate; oleic acid decyl ester, ethyl oleate, or C6-C12-alkanoic acid esters - especially caprylic/capric acid esters - with saturated fatty alcohols, especially C12-C18 saturated fatty alcohols.

Preferred is a mixture of paraffin - especially liquid paraffin - and C6-C12-alkanoic acid C12-C18-alkyl esters - especially caprylic/capric acid esters with C12-C18 saturated fatty alcohols (coco-caprylate/caprate, e.g. Cetiol®LC) as well as a mixture of isopropyl myristate and C6-C12-alkanoic acid C12-C18-alkyl esters (as just characterized above). The lipid(s) (f) are preferably present in a total amount of 3-7, especially 4-6, % of the total composition. If a mixture of paraffin and coco-caprylate/caprate is used, the amount of paraffin is preferably 1.5-3%, especially 2-2.8%, and the amount of coco-caprylate/caprate is preferably 1.5-3%, especially 2-2.8%, of the total composition.

(g) A non-ionic surfactant is e.g. an ester of a fatty acid, especially a C8-C18 fatty acid, with monohydroxy or, preferably, polyhydroxy compounds, e.g. ethylene glycol, glycerol, anhydrosorbitol or pentaerythritol. Another important group of non-ionic surfactants is characterized by compounds having at least one active hydrogen, e.g. fatty alcohols - especially a C8-C18 fatty alcohol -, fatty acids - especially a C8-C18 fatty acid -, sorbitan fatty acid esters, C1-C18-alkylphenols or C8-C18-alkylamines, that all are poly(oxyethylated), preferably with from 2 up to 40 ethylene glycol or ethylene oxide units.

Examples for the above mentioned non-ionic surfactants are partial glycerin fatty acid esters, such as glycerin monostearate; partial fatty acid esters of sorbitan or polyoxyethylene sorbitan, such as sorbitan monolaurate or polyethylene glycol (5 to 20) sorbitan monostearate or monooleate; polyoxyethylene (3 to 40) fatty alcohol ethers, such as polyoxyethylene (3 to 12) lauryl ethers or polyoxyethylene (5 to 40) cetostearyl ethers; polyoxyethylene fatty acid esters, such as polyoxyethylene (8 to 100) stearate; polyoxyethylene C4-C12-alkylphenyl ethers, e.g. polyoxyethylene (nonyl or octyl)phenyl ethers; or polyoxyethylene C8-C18-alkylamines, e.g. polyoxyethylene oleylamine. Preferred are polyoxyethylene (10 to 30) fatty alcohol ethers, in particular polyoxyethylene (20) cetostearyl ether (e.g. Cetomacrogol 1000). The non-ionic surfactant (g) is preferably present in an amount of from 1.5 up to 2.5% of the total composition.

(h) Ammonia is preferably applied in the form of a concentrated aqueous solution thereof. To adjust the pH of the total composition to 6.5-8 - preferably 7-8 -, said concentrated aqueous solution of ammonia is e.g. present in an amount of 0.2-3% (w/w), especially 0.5-2% (w/w), of the total composition. Sodium hydroxide (NaOH) and potassium hydroxide (KOH) are e.g. applied as concentrated, e.g. 30%, aqueous solutions thereof.

The compositions of the inventions may optionally include further routine excipients known in the art, for example perfumes and antimicrobial preservatives, e.g. benzyl alcohol, benzalkonium chloride or parabens (= C1-C7-alkyl esters of 4-hydroxybenzoic acid, e.g. methyl 4-hydroxybenzoate).

In the experimental emulsion-gel "B)" disclosed in US patent 4,917,886, the addition of a chemical stabilizer (sodium sulphite) was necessary to provide a chemically stable

formulation. In contrast thereto, the compositions of the present invention are chemically stable without a said stabilizer. In the absence of a stabilizer, however, e.g. the analytical work necessary to characterize the formulations and register them at the health authorities is simplified. Therefore, a further embodiment of the invention is characterized by the compositions disclosed hereinabove and hereinbefore, which compositions are devoid of a chemical stabilizer. Chemical stabilizers are e.g. antioxidants or chelating agents, e.g. EDTA.

The topical pharmaceutical compositions according to the invention are administered in a manner known per se. For example, they are applied e.g. once, twice or three times daily to the affected portions of the skin.

The invention further relates to a method of treating inflammatory diseases which comprises topically administering to a mammal in need of such treatment a therapeutically effective amount of one of the topical pharmaceutical compositions as specified herein above and below.

The manufacture of the topical pharmaceutical preparations is effected in a manner known per se, for example as described in the examples below.

The following examples are intended to illustrate the invention.

Example 1: An emulsion gel comprising 0.1% of diclofenac Na

<u>Ingredients</u>	<u>Amount (kg/100kg)</u>
(a) diclofenac sodium salt	0.10
(b) purified water	76.57
(c) isopropanol	10.0
(d) propylene glycol	5.0
(e) Carbomer 980	0.7
(f) paraffin, liquid	2.5
(f') coco-caprylate/caprate	2.5
(g) polyoxyethylene-20-cetostearyl ether	2.0
(h) ammonia, concentrated solution in water	<u>0.63</u>
	100.0

Manufacture: (a) is dissolved in (c), (d) and (h). Said solution is added to a mixture of (b) and (e). All remaining components – (f'), (f'') and (g) - are heated and slowly added to the former mixture. Upon mixing a homogeneous emulsion-gel is obtained.

Example 1a: An emulsion gel comprising 0.3% of diclofenac Na is obtained in an analogous manner to Example 1 but with using 0.30 of (a), 76.0 of (b), 0.9 of (e) and 0.8 of (h) instead.

Example 2: An emulsion gel comprising 0.1% of diclofenac Na [without (c)]

<u>Ingredients</u>	<u>Amount (kg/100kg)</u>
(a) diclofenac sodium salt	0.10
(b) purified water	74.9
(c) —	—
(d) propylene glycol	15.0
(e) Carbomer 974P	1.0
(f') paraffin, liquid	2.5
(f'') coco-caprylate/caprate	2.5
(g) polyoxyethylene-20-cetostearyl ether	2.0
(h) 30% aqueous NaOH solution	1.5
benzyl alcohol (as preservative)	<u>0.5</u>
	100.0

Example 3: A sprayable emulsion gel (being more fluid but still turbid and opaque) comprising 0.1% of diclofenac Na [without (c)]

<u>Ingredients</u>	<u>Amount (kg/100kg)</u>
(a) diclofenac sodium salt	0.10
(b) purified water	86.68
(c) —	—
(d) propylene glycol	5.0
(e) Carbomer 974P	0.3
(f') paraffin, liquid	2.5
(f'') coco-caprylate/caprate	2.5

(g) polyoxyethylene-20-cetostearyl ether	2.0
(h) 30% aqueous NaOH solution	0.42
<hr/>	
benzyl alcohol (as preservative)	<u>0.5</u>
	100.0

The emulsion-gel is manufactured in a manner analogous to Example 1.

Claims

1. A pharmaceutical composition intended for topical use, which composition is in the form of an opaque emulsion-gel and which composition comprises

- (a) 0.02-0.4% (w/w) of diclofenac sodium salt,
- (b) at least 50% (w/w) of water,
- (c) 0-30% (w/w) of at least one C2-C4-alkanol,
- (d) 3-20% (w/w) of a glycol solvent selected from the group consisting of propylene glycol and polyethylene glycol (200-1000),
- (e) 0.2-3% (w/w) of at least one gelling agent selected from the group consisting of carbomers,
- (f) 2-8% (w/w) of at least one lipid forming the oily phase of the emulsion-gel,
- (g) 1-5% (w/w) of at least one non-ionic surfactant, and
- (h) a basic agent selected from the group consisting of ammonia, sodium hydroxide and potassium hydroxide to adjust the pH of the total composition to 6.5-8.

2. A composition according to claim 1, which comprises

- (a) 0.05-0.3% of diclofenac sodium salt,
- (b) 60-92% of water,
- (c) 0-25% of ethanol, isopropanol, or mixtures thereof,
- (d) 3-20% of propylene glycol,
- (e) 0.3-2% of at least one gelling agent selected from the group consisting of carbomers,
- (f) 3-7% of at least one lipid forming the fatty phase of the emulsion-gel,
- (g) 1-3% of at least one non-ionic surfactant, and
- (h) ammonia to adjust the pH of the total composition to 6.5-8.

3. A composition according to claim 1 or claim 2, which comprises (c) in an amount of from 0 up to 5% and (d) in an amount of from 3 up to 20% of the total composition.

4. A composition according to claim 3, which is devoid of the component (c).

5. A composition according to any one of claims 1-4, which comprises as gelling agent (e) carbomer 980, carbomer 940, carbomer 981, carbomer 941, carbomer 974, carbomer 934, carbomer 910, or a mixture of any of said carbomers.

6. A composition according to any one of claims 1-5, which comprises as lipid (f) either a mixture of paraffin and C6-C12-alkanoic acid C12-C18-alkyl esters, or a mixture of isopropyl myristate and C6-C12-alkanoic acid C12-C18-alkyl esters.

7. A composition according to claim 3 or claim 4, which comprises as lipid (f) isopropyl myristate.

8. A composition according to any one of claims 1-7, which comprises as non-ionic surfactant (g) a polyoxyethylene (10 to 30) fatty alcohol ether.

9. A composition according to any one of claims 1-8, wherein the pH of the total composition is adjusted to pH 7-8.

10. A composition according to any one of claims 1-9, which compositions are devoid of a chemical stabilizer.

Topical Composition**Abstract of the Disclosure**

The invention relates to beneficial topical pharmaceutical compositions comprising the diclofenac sodium salt in unusually low amounts. Said compositions represent opaque emulsion-gels, in which diclofenac is kept fully dissolved.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.